

New York, 2006

Primary and secondary syphilis

- **CASES:** New York reported 736 cases in 2006.
- **RATES:** New York ranked 12 among 50 states, DC, and 3 territories with 3.8 cases per 100,000 population compared to the total rate of 3.3 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In New York, the rate among males was 7.6 per 100,000 population compared to the total male rate of 5.7 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the total female rate of 1.0 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


In New York, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 9.8 among blacks, 6.0 among Hispanics, 1.2 among Asian/Pacific Islanders, and 7.0 among American Indians/Alaska Natives.

- The rate among blacks was 4.9 times that of whites.
- The rate among Hispanics was 3 times that of whites.
- The rate among American Indians/Alaska Natives was 3.5 times that of whites.

Congenital syphilis

- **CASES:** New York reported 15 cases in 2006.
- **RATES:** New York ranked 18 among 26 states, DC, and 1 territory reporting congenital syphilis cases with a rate of 6.0 cases per 100,000 live births compared to the total rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, New York, 1997-2006



Total P&S Syphilis Rates by County, New York, 2006

